



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Limits Preview/Index History Clipboard Details

Abstract Text

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: J Allergy Clin Immunol. 2000 Feb;105(2 Pt 1):378-84.

[Related Articles.](#)

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

Mutational analysis of the IgE-binding epitopes of P34/Gly m Bd 30K.

Helm RM, Cockrell G, Connaughton C, West CM, Herman E, Sampson HA, Bannon GA, Burks AW.

Department of Pediatrics, University of Arkansas for Medical Sciences, Arkansas Children's Nutrition Center, Little Rock, AR 72202, USA.

BACKGROUND: Peanuts and soybeans are 2 foods that have been shown to be responsible for many atopic disorders. Because of their nutritional benefit, soybean proteins are now being used increasingly in a number of food products. Previous studies have documented multiple allergens in soybean extracts, including glycinin, beta-conglycinin, and the P34/Gly m Bd 30K protein. **OBJECTIVE:** Our overall goal was to identify soybean-specific allergens to begin to understand molecular and immunochemical characteristics of legume proteins. The specific aim of the current investigation was to identify the essential amino acid residues necessary for IgE binding in the 5 distinct immunodominant epitopes of P34/Gly m Bd 30K. **METHODS:** Serum IgE from 6 clinically sensitive soybean-allergic individuals was used to identify P34/Gly m Bd 30K in the native and single amino acid substituted peptides with use of the SPOTS peptide synthesis technique to determine critical amino acids required for IgE binding. **RESULTS:** The intensity of IgE binding and epitope recognition by serum IgE from the individuals varied substantially. With use of serum from 6 clinically soybean-sensitive individuals, 2 of the 5 immunodominant epitopes could be mutagenized to non-IgE binding peptide. **CONCLUSIONS:** Single-site amino acid substitution of the 5 immunodominant epitopes of Gly m Bd 30K with alanine revealed that IgE binding could be reduced or eliminated in epitopes 6 and 16 in the serum obtained from 6 soybean-sensitive patients.

Publication Types:

- Clinical Trial
- Controlled Clinical Trial

PMID: 10669862 [PubMed - indexed for MEDLINE]